

Identification of latent neosporosis in sheep in Tehran, Iran by polymerase chain reaction using primers specific for the *Nc-5* gene

**Authors:**

Mohsen Arbabi¹
Amir Abdoli¹
Abdolhossein Dalimi²
Majid Pirestani²

Affiliations:

¹Department of Parasitology,
Kashan University of Medical
Sciences, Iran

²Department of Parasitology,
Tarbiat Modares University,
Iran

Research Project no:

92019029

Corresponding author:

Amir Abdoli,
a.abdoli@modares.ac.ir

Dates:

Received: 10 Sept. 2015
Accepted: 24 Sept. 2015
Published: 11 Aug. 2016

How to cite this article:

Arbabi, M., Abdoli, A., Dalimi, A. & Pirestani, M., 2016, 'Identification of latent neosporosis in sheep in Tehran, Iran by polymerase chain reaction using primers specific for the *Nc-5* gene', *Onderstepoort Journal of Veterinary Research* 83(1), a1058. <http://dx.doi.org/10.4102/ojvr.v83i1.1058>

Copyright:

© 2016. The Authors.
Licensee: AOSIS. This work
is licensed under the
Creative Commons
Attribution License.

Little is known about latent infection and molecular characterisation of *Neospora caninum* in sheep (*Ovis aries*). In this study, 330 sheep samples (180 hearts and 150 brains) were analysed for *N. caninum* DNA by nested polymerase chain reaction (PCR) targeting the *Nc-5* gene. *Neospora caninum* DNA was detected in 3.9% (13/330) of sheep samples. The parasite's DNA was detected in 6.7% of heart samples (12/180) and 0.7% (1/150) of brain samples. No clinical signs were recorded from infected or uninfected animals. Sequencing of the genomic DNA revealed 96% – 99% similarity with each other and 95.15% – 100% similarity with *N. caninum* sequences deposited in GenBank. To our knowledge, this is the first report on the use of PCR to identify latent neosporosis in sheep in Iran. The results of this study have the potential to contribute to our understanding of the role of *N. caninum*-infected sheep in the epidemiology of neosporosis.

Introduction

Neospora caninum is a worldwide protozoan having a variety of animal hosts (Dubey & Schares 2011; Dubey, Schares & Ortega-Mora 2007). Domestic and wild canids are definitive, whereas different bird and mammalian species (such as cattle, water buffalo, and sheep) serve as intermediate hosts (Dubey & Schares 2011). Abortion, especially in dairy cattle, is one of the major consequences of neosporosis in animal husbandry (Almeria & López-Gatius 2013) that lead to significant economic losses (Reichel *et al.* 2013). Moreover, ovine abortion and reproductive failure due to neosporosis have been reported in several studies (Dubey & Lindsay 1990; Howe *et al.* 2008, 2012; Jolley *et al.* 1999; Moreno *et al.* 2012; Pena *et al.* 2007). In different studies, antibodies to *N. caninum* have been detected in 1.1% – 8.3% of sheep in the west of Iran (Ezatpour *et al.* 2015; Gharekhani & Heidari 2014), 27.7% in Pakistan (Nasir *et al.* 2012), 2.1% in Turkey (Gökçe *et al.* 2015), 10.3% in China (Liu *et al.* 2015), 16.8% in Greece (Anastasia *et al.* 2013), 3% in Argentina (Hecker *et al.* 2013), and 13.1% in south-eastern Brazil (Da Silva Andrade *et al.* 2012). However, there is little information describing the detection of nucleic acids resulting from latent neosporosis in sheep.

Until now, different genes such as internal transcribed spacer sequences, 18S-like ribosomal DNA (small-subunit rDNA), and *Nc-5* genes have been used for molecular diagnosis of neosporosis (reviewed by Al-Qassab, Reichel & Ellis [2010]; Goodswen, Kennedy & Ellis [2013]). However, studies have indicated that the *Nc-5* gene is one of the most highly sensitive and specific for the detection of neosporosis (Almeria *et al.* 2002; Dubey *et al.* 2014; Hughes *et al.* 2006; Kaufmann *et al.* 1996; Paula *et al.* 2004; Yamage, Flechtner & Gottstein 1996) because it is repeated in the *N. caninum* sequence (Al-Qassab *et al.* 2010). Hence, the main objective of this study was to investigate detection and molecular characterisation of latent neosporosis in sheep (*Ovis aries*) in Tehran, Iran, by polymerase chain reaction (PCR) using primers specific for the *Nc-5* gene.

Materials and methods

Animals and study area

A total of 330 samples from healthy slaughtered sheep (180 hearts and 150 brains) were purchased from an abattoir in Vavan (located in the vicinity of Tehran) from April to September 2014. The animals tested originated from different counties (Eslamshahr, Shahriar, Robatkarim), all of which are located between 50 km and 200 km from Tehran. These locations have hot summers and moderate winters. No clinical signs such as fever, lymphadenitis, nasal and ocular discharges, or jaundice were recorded in any of the animals before slaughter.

Read online:

Scan this QR
code with your
smart phone or
mobile device
to read online.

DNA extraction

The whole brain and heart of each sheep were individually rinsed with distilled water, packaged, and refrigerated. Approximately 200 g – 250 g of different segments of brain and heart were homogenised with a pestle and mortar in liquid nitrogen, and DNA was extracted using a phenol–chloroform extraction method as described in our recent report (Abdoli *et al.* 2015). To prevent DNA cross-contamination, all materials that were used between different tissue samples were decontaminated with sodium hypochlorite solution (2.5%) and rinsed with distilled water. The concentration of DNA was determined by NanoDrop spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA) for each sample. Overall, the DNA concentration ranged between 150 ng/μL and 200 ng/μL.

Nested polymerase chain reaction

Nested PCR was conducted using specific primers for the *Nc-5* gene. The first round of PCR was conducted using a pair of *N. caninum*-specific primers, Np21plus (5'-CCCAGTGGTCCAATCCTGTAAAC-3') and Np6plus (5'-CTCGCCAGTCCAACCTACGTCTTCT-3') (Muller *et al.* 1996). Nested PCR was performed with the primers Np6 (5'-CAGTCAACCTACGTCTTCT-3') and Np7 (5'-GGGTGAACCGAGGGAGTTG-3') (Hughes *et al.* 2006). Each amplification was performed in 20-μL reaction mixtures containing 10 μL of 2x master mixes (DFS Master Mix, BIORON GmbH, Ludwigshafen, Germany), each of the respective primers (10 pmol for the first round reaction and 25 pmol for nested PCR), 7 μL of distilled water, and 1 μL of template DNA. One microlitre of the first round product was used as the template for nested PCR. For each reaction, a negative control (double distilled water) and a positive control (DNA extracted from the *Nc-5* strain of *N. caninum*) were included. Amplification was performed with initial denaturation for 5 minutes at 94 °C, followed by 40 cycles at 94 °C for 40 seconds (denaturation), annealing at 62 °C in the first round, and 56 °C in nested PCR for 40 seconds, extension at 72 °C for 40 seconds, and final extension at 72 °C for 10 minutes. PCR products were electrophoresed on a 1.5% agarose gel stained with safe stain (Sinaclon, Tehran, Iran) and visualised under ultraviolet trans-illumination.

Nucleotide sequence analysis

Four positive PCR products (from the second reaction) were amplified with a master mix containing *Pfu* DNA polymerase (Thermo Fisher Scientific, Waltham, USA, cat. no. EP0501), extracted from the gel (Vivantis gel purification kit, Selangor Darul Ehsan, Malaysia) according to the manufacturer's protocols. Then the products were sequenced in the forward and reverse directions by Sequetech (Mountain View, CA, USA) (Abdoli *et al.* 2015). The sequences were edited with BioEdit sequence alignment editor (Hall 1999), aligned with *Nc-5* partial sequences from other hosts by ClustalX2.12 (Larkin *et al.* 2007) and compared with sequences of *N. caninum* available in GenBank. Phylogenetic trees were


inferred and evolutionary analyses were conducted using the Tamura three-parameter option of the neighbour-joining model with MEGA6 software (<http://www.megasoftware.net/>) (Tamura *et al.* 2013). The bootstrap scores were calculated for 1000 replicates (Tamura *et al.* 2013).

Results

Neospora caninum DNA was detected in 13 out of 330 sheep samples (3.9%). The infection rates in the heart and brain samples were 6.7% (12/180) and 0.7% (1/150), respectively. Four nucleotide sequences of the *Nc-5* gene with a length of 227 bp (Figure 1) were submitted to the GenBank database (GenBank accession numbers KR106181, KR106182, KR106183, KR106184). The results demonstrated our sequences shared 96% – 99% similarity with each other (Figures 2 and 3) and 95% – 100% similarity with *N. caninum* deposited in GenBank (Appendix Figure 1). Phylogenetic trees showed intraspecific variations between our isolates and other *N. caninum* specimens deposited in GenBank (Figure 2). Analysis of our sequences showed 96.9% – 97.8% similarity with *N. caninum* isolated from sheep (DQ077661) in the UK and 96.9% – 99.1% similarity with *N. caninum* isolated from sparrows (*Passer domesticus*) in Iran. Interestingly, one of our samples (KR106181) showed 100% similarity with *N. caninum* isolated from wolves (*Canis lupus*) (KF649846) in the United States.

Discussion

Although an association between ovine abortion and neosporosis has been reported in different studies (Dubey & Lindsay 1990; Howe *et al.* 2008, 2012; Jolley *et al.* 1999; Moreno *et al.* 2012; Pena *et al.* 2007), there is little information describing molecular detection of latent neosporosis in sheep. Here, we found a total infection rate of 3.9% (13/330) in our sheep samples. Interestingly, 12 out of 13 positive samples were detected in the hearts and one positive sample was diagnosed in the brain. In previous studies, the seroprevalence of *N. caninum* has been reported in a range of 1.1% – 8.3% of sheep from the west of Iran (Ezatpour *et al.* 2015; Gharekhani & Heidari 2014). Moreover, *N. caninum* DNA was detected in 8.5% (Asadpour *et al.* 2013) and 0.9% of aborted ovine fetuses



M, 100 bp DNA marker; Lane 1, positive control; Lane 2, negative control; Lanes 3, 4, 5, 7, positive samples; Lane 6, negative samples.

FIGURE 1: Polymerase chain reaction products of four *Neospora caninum* positive samples.

neosporosis in experimentally infected mice. Among 19 combinations of forward and reverse primers, the Np21/Np6, Np7/Np6, and Np21/Np4 primer pairs were able to detect at least 10 pg genomic DNA with a specific single band (Yamage *et al.* 1996). The *Nc-5* gene can also discriminate *N. caninum* from other related apicomplexan parasites (*Toxoplasma gondii* and *Sarcocystis* species) (Kaufmann *et al.* 1996). Thus, the *Nc-5* gene has been used as a highly sensitive and specific gene for detection of neosporosis (Almeria *et al.* 2002; Dubey *et al.* 2014; Hughes *et al.* 2006; Paula *et al.* 2004; Yamage *et al.* 1996). Hence, we selected the *Nc-5* gene for sensitive and specific detection of neosporosis in the current study.

We also sequenced four positive samples for phylogenetic analysis. We found that our sequences displayed similarity levels of 96% – 99% with each other (Figure 3) and 95% – 100% with *N. caninum* sequences deposited in GenBank (Appendix Figure 1). In comparison with molecular diagnosis, few studies have been conducted on the phylogenetic analysis of *N. caninum* with the *Nc-5* gene (Auriemma *et al.* 2014; Čobádiová *et al.* 2013; Hughes *et al.* 2006). BLAST analyses indicated greater than 94% (Čobádiová *et al.* 2013), 96% (Auriemma *et al.* 2014), and 97% (Hughes *et al.* 2006) similarities between their sequences and other *N. caninum* sequences deposited in GenBank. It therefore seems that the *Nc-5* gene is not a suitable biomarker for phylogenetic analysis and discrimination of genetic diversity for *N. caninum*. Instead, this gene is rather a highly sensitive and specific biomarker for the diagnosis of neosporosis. The use of ribosomal DNA, ITS-1, and recently microsatellites have been recommended for discriminating between *N. caninum* isolates (Al-Qassab *et al.* 2010).

Taken together, the results of this study provide molecular and epidemiological information about latent *N. caninum* infection in sheep in Iran. It can be expected that in future these results will contribute to revealing the role of latent *N. caninum* infection in the epidemiology of neosporosis in sheep.

Acknowledgements

This work was supported by the Iran National Science Foundation (grant no. 92019029). The authors would like to thank Mr. Mahdi Rezapour, Ali Karaji, and Ali Shafiei for their help in preparing the samples.

Competing interests

The authors declare that they have no financial or personal relationships that may have inappropriately influenced them in writing this article.

Authors' contributions

M.A. conceived the study design, analysed and interpreted the data and reviewed the manuscript; A.A. collected the data, prepared the samples, performed molecular assays, designed the tables and figures and wrote the manuscript. M.P. performed the molecular assays and sequencing analysis; A.D. conceived the study design and analysed and interpreted the data.

References

- Abdoli, A., Arbabi, M., Dalimi, A. & Pirestani, M., 2015, 'Molecular detection of *Neospora caninum* in house sparrow (*Passer domesticus*) in Iran', *Avian Pathology* 44(2), 319–322. <http://dx.doi.org/10.1080/03079457.2015.1050583>
- Almeria, S., Ferrer, D., Pabón, M., Castella, J. & Manas, S., 2002, 'Red Foxes (*Vulpes vulpes*) are a natural intermediate host of *Neospora caninum*', *Veterinary Parasitology* 107, 287–294. [http://dx.doi.org/10.1016/S0304-4017\(02\)00162-0](http://dx.doi.org/10.1016/S0304-4017(02)00162-0)
- Almeria, S. & López-Gatius, F., 2013, 'Bovine neosporosis: Clinical and practical aspects', *Research in Veterinary Science* 95, 303–309. <http://dx.doi.org/10.1016/j.rvsc.2013.04.008>
- Al-Qassab, S.E., Reichel, M.P. & Ellis, J.T., 2010, 'On the biological and genetic diversity in *Neospora caninum*', *Diversity* 2, 411–438. <http://dx.doi.org/10.3390/d2030411>
- Anastasia, D., Elias, P., Nikolaos, P., Charilaos, K. & Nektarios, G., 2013, 'Toxoplasma gondii and Neospora caninum seroprevalence in dairy sheep and goats mixed stock farming', *Veterinary Parasitology* 198, 387–390. <http://dx.doi.org/10.1016/j.vetpar.2013.09.017>
- Andrianarivo, A.G., Anderson, M.L., Rowe, J.D., Gardner, I.A., Reynolds, J.P., Choromanski, L. *et al.*, 2005, 'Immune responses during pregnancy in heifers naturally infected with *Neospora caninum* with and without immunization', *Parasitology Research* 96, 24–31. <http://dx.doi.org/10.1007/s00436-005-1313-y>
- Asadpour, R., Jafari-Joozani, R. & Salehi, N., 2013, 'Detection of *Neospora caninum* in ovine abortion in Iran', *Journal of Parasitic Diseases* 37, 105–109.
- Auriemma, C., Lucibelli, M.G., Borriello, G., De Carlo, E., Martucciello, A., Schiavo, L. *et al.*, 2014, 'PCR detection of *Neospora caninum* in water Buffalo Foetal Tissues', *Acta Parasitologica* 59, 1–4. <http://dx.doi.org/10.2478/s11686-014-0201-y>
- Bishop, S., King, J., Windsor, P., Reichel, M.P., Ellis, J. & Šlapeta, J., 2010, 'The first report of ovine cerebral neosporosis and evaluation of *Neospora caninum* prevalence in sheep in New South Wales', *Veterinary Parasitology* 170, 137–142. <http://dx.doi.org/10.1016/j.vetpar.2010.01.030>
- Čobádiová, A., Vichova, B., Majlathova, V. & Reiterová, K., 2013, 'First molecular detection of *Neospora caninum* in European Brown Bear (*Ursus arctos*)', *Veterinary Parasitology* 197, 346–349. <http://dx.doi.org/10.1016/j.vetpar.2013.05.005>
- Da Silva Andrade, G., Bruhn, F.R.P., Rocha, C.M.B.M., De Sá Guimarães, A., Gouveia, A.M.G. & Guimarães, A.M., 2012, 'Seroprevalence and risk factors for *Neospora caninum* in sheep in The State Minas Gerais, Southeastern Brazil', *Veterinary Parasitology* 188, 168–171. <http://dx.doi.org/10.1016/j.vetpar.2012.03.006>
- Dubey, J., Jenkins, M., Ferreira, L., Choudhary, S., Verma, S., Kwok, O. *et al.*, 2014, 'Isolation of viable *Neospora caninum* from brains of Wild Gray Wolves (*Canis lupus*)', *Veterinary Parasitology* 201, 150–153. <http://dx.doi.org/10.1016/j.vetpar.2013.12.032>
- Dubey, J. & Lindsay, D.S., 1990, '*Neospora caninum* induced abortion in sheep', *Journal of Veterinary Diagnostic Investigation* 2, 230–233. <http://dx.doi.org/10.1177/104063879000200316>
- Dubey, J. & Schares, G., 2011, 'Neosporosis in animals – The last five years', *Veterinary Parasitology* 180, 90–108. <http://dx.doi.org/10.1016/j.vetpar.2011.05.031>
- Dubey, J., Schares, G. & Ortega-Mora, L., 2007, 'Epidemiology and control of Neosporosis and *Neospora caninum*', *Clinical Microbiology Reviews* 20, 323–367. <http://dx.doi.org/10.1128/CMR.00031-06>
- Ezatpour, B., Alirezai, M., Hassanvand, A., Zibaei, M., Azadpour, M. & Ebrahimzadeh, F., 2015, 'The first report of *Neospora caninum* prevalence in aborted and healthy sheep from west of Iran', *Comparative Clinical Pathology* 24, 19–22. <http://dx.doi.org/10.1007/s00580-013-1846-x>
- Gharekhani, J. & Heidari, H., 2014, 'Serology based comprehensive study of *Neospora* infection in domestic animals in Hamedan Province, Iran', *Journal Advances Veterinary Animal Research* 1, 119–124. <http://dx.doi.org/10.5455/javar.2014.a23>
- Gökçe, G., Mor, N., Kirmizigül, A., Bozukluhan, K. & Erkilic, E., 2015, 'The first report of seropositivity for *Neospora caninum* in sheep from Turkey', *Israel Journal of Veterinary Medicine* 70, 40–44.
- Goodswen, S.J., Kennedy, P.J. & Ellis, J.T., 2013, 'A review of the infection, genetics, and evolution of *Neospora caninum*: From the past to the present', *Infection, Genetics and Evolution* 13, 133–150. <http://dx.doi.org/10.1016/j.meegid.2012.08.012>
- Hall, T.A., 1999, *Bioedit: A user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT*, Nucleic Acids Symposium Series, pp. 95–98, Oxford University Press, Oxford.
- Hecker, Y.P., Moore, D.P., Manazza, J.A., Unzaga, J.M., Späth, E.J., Pardini, L.L. *et al.*, 2013, 'First report of seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in dairy sheep from Humid Pampa, Argentina', *Tropical Animal Health and Production* 45, 1645–1647. <http://dx.doi.org/10.1007/s11250-013-0396-1>
- Hemphill, A., Vonlaufen, N. & Naguleswaran, A., 2006, 'Cellular and immunological basis of the host-parasite relationship during infection with *Neospora caninum*', *Parasitology* 133, 261–278. <http://dx.doi.org/10.1017/S0031182006000485>
- Howe, L., Collett, M., Pattison, R., Marshall, J., West, D. & Pomroy, W., 2012, 'Potential involvement of *Neospora caninum* in naturally occurring ovine abortions in New Zealand', *Veterinary Parasitology* 185, 64–71. <http://dx.doi.org/10.1016/j.vetpar.2011.10.033>
- Howe, L., West, D., Collett, M., Tattersfield, G., Pattison, R., Pomroy, W. *et al.*, 2008, 'The role of *Neospora caninum* in three cases of unexplained ewe abortions in the Southern North Island of New Zealand', *Small Ruminant Research* 75, 115–122. <http://dx.doi.org/10.1016/j.smallrumres.2007.08.001>
- Hughes, J., Williams, R., Morley, E., Cook, D., Terry, R., Murphy, R. *et al.*, 2006, 'The prevalence of *Neospora caninum* and co-infection with *Toxoplasma gondii* by PCR analysis in naturally occurring mammal populations', *Parasitology* 132, 29–36. <http://dx.doi.org/10.1017/S0031182005008784>

- Jolley, W., Mcallister, M., Mcguire, A. & Wills, R., 1999, 'Repetitive abortion in *Neospora*-infected ewes', *Veterinary Parasitology* 82, 251–257. [http://dx.doi.org/10.1016/S0304-4017\(99\)00017-5](http://dx.doi.org/10.1016/S0304-4017(99)00017-5)
- Kaufmann, H., Yamage, M., Roditi, I., Dobbelaere, D., Dubey, J., Holmdahl, O. *et al.*, 1996, 'Discrimination of *Neospora caninum* from *Toxoplasma gondii* and other apicomplexan parasites by hybridization and PCR', *Molecular and Cellular Probes* 10, 289–297. <http://dx.doi.org/10.1006/mcpr.1996.0038>
- Larkin, M.A., Blackshields, G., Brown, N., Chenna, R., Mcgettigan, P.A., Mcwilliam, H. *et al.*, 2007, 'Clustal W and clustal X version 2.0', *Bioinformatics* 23, 2947–2948. <http://dx.doi.org/10.1093/bioinformatics/btm404>
- Liu, Z.-K., Li, J.-Y. & Pan, H., 2015, 'Seroprevalence and risk factors of *Toxoplasma gondii* and *Neospora caninum* infections in small ruminants in China', *Preventive Veterinary Medicine* 118, 488–492. <http://dx.doi.org/10.1016/j.prevetmed.2014.12.017>
- Magaña, A., Sánchez, F., Villa, K., Rivera, L. & Morales, E., 2015, 'Systemic neosporosis in a dog treated for immune-mediated thrombocytopenia and hemolytic anemia', *Veterinary Clinical Pathology* 44, 592–596. <http://dx.doi.org/10.1111/vcp.12287>
- Mazuz, M.L., Shkap, V., Wollkomirsky, R., Leibovich, B., Savitsky, I., Fleiderovitz, L. *et al.*, 2016, '*Neospora caninum*: Chronic and congenital infection in consecutive pregnancies of mice', *Veterinary Parasitology* 219, 66–70. <http://dx.doi.org/10.1016/j.vetpar.2016.01.013>
- Moreno, B., Collantes-Fernández, E., Villa, A., Navarro, A., Regidor-Cerrillo, J. & Ortega-Mora, L., 2012, 'Occurrence of *Neospora caninum* and *Toxoplasma gondii* infections in ovine and caprine abortions', *Veterinary Parasitology* 187, 312–318. <http://dx.doi.org/10.1016/j.vetpar.2011.12.034>
- Muller, N., Zimmermann, V., Hentrich, B. & Gottstein, B., 1996, 'Diagnosis of *Neospora caninum* and *Toxoplasma gondii* infection by PCR and DNA hybridization immunoassay', *Journal Clinical Microbiology* 34, 2850–2852.
- Nasir, A., Ashraf, M., Khan, M., Javeed, A., Yaqub, T., Avais, M. *et al.*, 2012, 'Prevalence of *Neospora caninum* antibodies in sheep and goats in Pakistan', *Journal Parasitology* 98, 213–215. <http://dx.doi.org/10.1645/GE-2863.1>
- Şuteu, O., Paştiu, A., Györke, A., Avram, E. & Cozma, V., 2013, 'Molecular detection of *Neospora caninum* in slaughtered goat kids from Romania', *Science Parasitology* 14, 43–46.
- Pabón, M., López-Gatius, F., García-Ispuerto, I., Bech-Sabat, G., Nogareda, C. & Almería, S., 2007, 'Chronic *Neospora caninum* infection and repeat abortion in dairy cows: A 3-year study', *Veterinary Parasitology* 147, 40–46. <http://dx.doi.org/10.1016/j.vetpar.2007.03.017>
- Paula, V., Rodrigues, A., Richtzenhain, L., Cortez, A., Soares, R. & Gennari, S., 2004, 'Evaluation of a PCR based on primers to *Nc5* gene for the detection of *Neospora caninum* in brain tissues of bovine aborted fetuses', *Veterinary Research Communications* 28, 581–585. <http://dx.doi.org/10.1023/B:VERC.0000042877.07684.89>
- Pena, H., Soares, R., Ragozo, A., Monteiro, R., Yai, L., Nishi, S. *et al.*, 2007, 'Isolation and molecular detection of *Neospora caninum* from naturally infected sheep from Brazil', *Veterinary Parasitology* 147, 61–66. <http://dx.doi.org/10.1016/j.vetpar.2007.03.002>
- Reichel, M.P., Alejandra Ayanegui-Alcerreca, M., Gondim, L.F. & Ellis, J.T., 2013, 'What is the global economic impact of *Neospora caninum* in cattle—the billion dollar question', *International Journal for Parasitology* 43, 133–142. <http://dx.doi.org/10.1016/j.ijpara.2012.10.022>
- Rettigner, C., De Meerschman, F., Focant, C., Vanderplasschen, A. & Losson, B., 2004, 'The vertical transmission following the reactivation of a *Neospora caninum* chronic infection does not seem to be due to an alteration of the systemic immune response in pregnant cba/ca mice', *Parasitology* 128, 149–160. <http://dx.doi.org/10.1017/S0031182003004402>
- Santos, S.L., De Souza Costa, K., Gondim, L.Q., Da Silva, M.S.A., Uzeda, R.S., Abe-Sandes, K. & Gondim, L.F.P., 2010, 'Investigation of *Neospora caninum*, *Hammondia* sp., and *Toxoplasma gondii* in tissues from slaughtered beef cattle in Bahia, Brazil', *Parasitology Research* 106, 457–461. <http://dx.doi.org/10.1007/s00436-009-1686-4>
- Sasani, F., Javanbakht, J., Seifori, P., Fathi, S. & Hassan, M.A., 2013, '*Neospora caninum* as causative agent of ovine encephalitis in Iran', *Pathology Discovery* 1, 5. <http://dx.doi.org/10.7243/2052-7896-1-5>
- Silva, M.S., Uzeda, R.S., Costa, K.S., Santos, S.L., Macedo, A.C., Abe-Sandes, K. & Gondim, L.F.P., 2009, 'Detection of *Hammondia Heydorni* and related coccidia (*Neospora caninum* and *Toxoplasma gondii*) in goats slaughtered in Bahia, Brazil', *Veterinary Parasitology* 162, 156–159. <http://dx.doi.org/10.1016/j.vetpar.2009.02.007>
- Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S., 2013, 'Mega6: Molecular evolutionary genetics analysis version 6.0', *Molecular Biology and Evolution* 30, 2725–2729. <http://dx.doi.org/10.1093/molbev/mst197>
- Yamage, M., Flechtner, O. & Gottstein, B., 1996, '*Neospora caninum*: Specific oligonucleotide primers for the detection of Brain "Cyst" dna of experimentally infected nude mice by the polymerase chain reaction (PCR)', *Journal Parasitology* 82, 272–279. <http://dx.doi.org/10.2307/3284160>

Appendix starts on the next page →

Appendix 1

| | | |
|----------|--|-----|
| EF581827 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| GU194965 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KF649844 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| EU073599 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| kp702735 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KR106185 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| kp702736 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| DQ077662 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| HM031966 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KR106182 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KR106183 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| DQ077661 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KF649848 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| kp702737 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| DQ132438 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KR106181 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KF649846 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| EF202080 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| GU194963 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| GU194964 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| GU194961 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| GU194962 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KR106184 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| DQ077663 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| DQ132437 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| EU073600 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| KF649847 | CAGTCAACCTACGTCTTCTGCCCTTCCCTCGTCCGCTTGCTCCCTATGCATAAATCTCCC | 60 |
| | *****.***.*****.***.*****. | |
| EF581827 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| GU194965 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KF649844 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| EU073599 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| kp702735 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KR106185 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 119 |
| kp702736 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 119 |
| DQ077662 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| HM031966 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KR106182 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KR106183 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| DQ077661 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KF649848 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| kp702737 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| DQ132438 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KR106181 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KF649846 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| EF202080 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| GU194963 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| GU194964 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| GU194961 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| GU194962 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KR106184 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| DQ077663 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| DQ132437 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| EU073600 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| KF649847 | CCGTCATCAGTGCCGCGCGGTGTTGCCTCAACACAGAACTGAACCTCTGGATAAGTATCA | 120 |
| | ..*.*****.*****.*****.*****.*****.*.*****.* | |

Our sheep samples are represented in red (accession nos. KR106181, KR106182, KR106183, KR106184). An asterisk represents an exact match between all sequences; a dot (•) represents a mismatch in at least one sequence.

FIGURE 1-A1: Sequence alignment of *Nc-5* gene of *Neospora caninum* from mammalian and bird hosts.

