ISSN: (Online) 2219-0635, (Print) 0030-2465

Page 1 of 1

Corrigendum: Anthelmintic activity of acetone extracts from South African plants used on egg hatching of *Haemonchus contortus*



Authors:

Gerda Fouche¹ ⁽²⁾ Bellonah M. Sakong² ⁽²⁾ Olubukola T. Adenubi² ⁽²⁾ Elizabeth Pauw² Tlabo Leboho¹ ⁽²⁾ Mbokota C. Khosa³ ⁽³⁾ Kevin W. Wellington¹ ⁽³⁾ Jacobus N. Eloff² ⁽³⁾

Affiliations:

¹CSIR Biosciences, Pretoria, South Africa

²Department of Paraclinical Sciences, University of Pretoria, South Africa

³Agricultural Research Council – Tropical and Subtropical Crops, Nelspruit, South Africa

Corresponding author: Kevin Wellington, kwellington@csir.co.za

Dates: Published: 27 July 2021

How to cite this article: Fouche, G., Sakong, B.M., Adenubi, O.T., Pauw, E., Leboho, T., Khosa, M.C. et al., 2021, 'Corrigendum: Anthelmintic activity of acetone extracts from South African plants used on egg hatching of *Haemonchus contortus'*, *Onderstepoort Journal of Veterinary Research* 88(1), a1949. https://doi. org/10.4102/ojvr.v88i1.1949

Copyright: © 2021. The Authors. Licensee: AOSIS. This work is licensed under the Creative Commons Attribution License.





Scan this QR code with your smart phone or mobile device to read online. In the published version of this article, Fouche, G., Sakong, B.M., Adenubi, O.T., Pauw, E., Leboho, T., Wellington, K.W. et al., 2016, 'Anthelmintic activity of acetone extracts from South African plants used on egg hatching of *Haemonchus contortus*', *Onderstepoort Journal of Veterinary Research* 83(1), a1164. https://doi.org/10.4102/ojvr.v83i1.1164, the sixth author, Mbokota C. Khosa, was omitted from the 'Authors' and 'Affiliations' sections. The indicated author should be added as the sixth author, and the following affiliation should be added as his affiliation: Agricultural Research Council – Tropical and Subtropical Crops, Nelspruit, South Africa.

The Authors' contributions section is hereby update to:

Authors' contributions

G.F. conceptualised the study. M.C.K. was involved in the collection of some of the plant material and in the preparation of the extracts used in the biological screening assays. G.F., K.W.W. and T.L. carried out the literature search and plant selection. T.L. prepared the plant extracts. J.N.E. and E.P. arranged for sheep to be infected, collected the eggs and guided the study. B.M.S. performed the egg hatch assay on the extracts. M.C.K. was also involved in the fractionation and isolation process in the natural product chemistry laboratory. O.T.A. screened the extracts for toxicity on Vero cells. K.W.W. wrote the first draft of the manuscript.

This correction does not alter the study's findings of significance or the overall interpretation of the study results. The authors apologise for any inconvenience caused.

ISSN: (Online) 2219-0635, (Print) 0030-2465

Page 1 of 7

Anthelmintic activity of acetone extracts from South African plants used on egg hatching of *Haemonchus contortus*



Authors:

Gerda Fouche¹ Bellonah M. Sakong² Olubukola T. Adenubi² Elizabeth Pauw² Tlabo Leboho¹ Kevin W. Wellington¹ Jacobus N. Eloff²

Affiliations: ¹CSIR Biosciences, Pretoria, South Africa

²Department of Paraclinical Sciences, University of Pretoria, South Africa

Corresponding author: Kevin Wellington, kwellington@csir.co.za

Dates:

Received: 02 Feb. 2016 Accepted: 12 May 2016 Published: 29 July 2016

How to cite this article:

Fouche, G., Sakong, B.M., Adenubi, O.T., Pauw, E., Leboho, T., Wellington, K.W. *et al.*, 2016, 'Anthelmintic activity of acetone extracts from South African plants used on egg hatching of *Haemonchus contortus'*, *Onderstepoort Journal of Veterinary Research* 83(1), a1164. http://dx.doi. org/10.4102/ojvr.v83i1.1164

Copyright:

© 2016. The Authors. Licensee: AOSIS. This work is licensed under the Creative Commons Attribution License.

Read online:



Scan this QR code with your smart phone or mobile device to read online. The nematode, *Haemonchus contortus*, is responsible for major economic losses in the livestock industry. The management of parasites such as *H. contortus* has been through the use of synthetic parasiticides. This has resulted in the presence of residues in meat and milk, which affects food safety. The development of resistance to available anthelmintics coupled with their high cost has further complicated matters. This has led to the investigation of alternative methods to manage nematodes, including the use of plants and plant extracts as a potential source of novel anthelmintic. Acetone extracts were prepared from 15 South African plant species and their anthelmintic activity determined using the egg hatch assay (EHA). The leaf extract of *Cleome gynandra* had the best inhibitory activity ($68\% \pm 3\%$) at a concentration of 2.5 mg/mL, followed by the stem extract of *Maerua angolensis* ($65\% \pm 5\%$). The extracts had a relatively low toxicity on Vero cells determined by the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) cellular assay.

Introduction

Livestock production in tropical and developing countries is severely hampered by gastrointestinal parasites (Adejinmi & Harrison 1997; Hounzangbe-Adote *et al.* 2005). The gastrointestinal parasitic nematode, *Haemonchus contortus*, also known as the barber pole worm, resides in the gut of sheep and other livestock. It accounts for about 80% of the global parasite afflictions of diseased animals (Arosemena *et al.* 1999) and is notorious for its high pathogenicity (Angulo-Cubillán *et al.* 2010).

In small ruminants, gastrointestinal nematodes have traditionally been managed by the use of synthetic anthelmintic compounds (Mendoza de Gives *et al.* 1998). The systematic application of anthelmintic drugs, in an effort to manage infections produced by *H. contortus*, has led to the emergence of resistant strains (Akhtar *et al.* 2000; Prichard 1994). There have been reports of parasite resistance to anthelmintic drugs in many countries (Melo, Bevilaqua & Reis 2009; Schnyder *et al.* 2005), and multiple anthelmintic resistance has reached extreme levels (Torres-Acosta *et al.* 2012).

Certain drugs may also cause problems such as food residues and environmental pollution (Hammond, Fielding & Bishop 1997). This global problem has caused severe losses in productivity and is also the main restricting factor for the livestock sector (Melo *et al.* 2009; Waller 1994). Novel alternative methods are thus required.

Research has been conducted on plant species as alternative anthelmintics to manage gastrointestinal infections in small ruminants (Adamu, Naidoo & Eloff 2013; Batista *et al.* 1999; Slomp *et al.* 2009). The use of anthelmintic plant extracts may be sustainable and environmentally acceptable and could provide an alternative to synthetic anthelmintics. Furthermore, anthelmintic plant extracts have a mixture of active principles that could act in synergy, yielding the anthelmintic effect and limit the development of resistance. This differs from commercial drugs, which usually have only one molecule acting on the parasite when not administered as a combination formulation. Resistance is therefore likely to develop more slowly in the natural product.

The aim of this study was to determine the *in vitro* anthelmintic action of acetone extracts from 15 South African plant species used traditionally to control parasites such as *H. contortus* using the egg hatch assay (EHA). We used only one concentration as a first step in selecting plant species for in-depth follow-up research. The toxicity of the acetone extracts was also determined against Vero cells.

Materials and methods Plant material collection

Fifteen plants [*Aloe rupestris* Baker, *Antizoma angustifolia* (Burch.) Miers ex Harv., *Calpurnia aurea* ssp. *aurea* (Aiton) Benth., *Senna italica* subsp. *arachoides* (Burch.) Lock, *Cissus quadrangularis* L., *Clematis brachiata* Thunb., *Cleome gynandra* L., *Ficus sycomorus* L., *Hypoxis rigidula* Baker var rigidula, *Maerua angolensis* DC, *Monsonia angustifolia* E. Mey. ex A. Rich., *Pelargonium luridium* (Andrews) Sweet, *Schkuhria pinnata* (Lam.) Kuntze ex Thell., *Sclerocarya birrea* (A. Rich.) Hochst and *Tabernaemontana elegans* Stapf.] were selected on the basis of available literature and ethno-veterinary usage over many years at Council for Scientific and Industrial Research (CSIR) (unpublished data). These plants were collected from different locations in South Africa during the summer season.

Production of dried, ground plant material

Plant material was dried in an oven at 30 $^{\circ}$ C – 60 $^{\circ}$ C followed by grinding to fine particles using a hammer mill.

Preparation of the acetone extracts

The acetone extract was prepared by adding 200 mL of acetone to 20 g of each ground plant material that was stirred for 1 h. The extract was decanted and filtered, and the residue was re-extracted with the same volume of acetone once again for 1 h; the third time, the same volume of acetone was used but the mixture was stirred overnight. The extracts were combined and the acetone evaporated on a rotary evaporator. The yield that was obtained for each of the plant species is shown in Table 1.

Egg recovery and preparation

The method used was based on the World Association for the Advancement of Veterinary Parasitology (WAAVP) guidelines described by Adamu *et al.* (2013). Briefly, *H. contortus* eggs were collected from sheep that were housed indoors on a concrete floor. Approximately 10 g – 15 g of sheep faecal pellets were crushed in water to form a slurry and cleared of organic debris by serially filtering it through sieves of pore sizes 150 μ m, 63 μ m and 20 μ m. The eggs were collected on a 20- μ m sieve and washed off with a 40% sugar solution (density 1.18) into 50-mL centrifuge tubes. The tubes were then centrifuged for 5 min at 1000 rpm to separate the floating eggs from other debris. The supernatant was decanted on a 20- μ m sieve, and the eggs were washed off with water and collected in a 500-mL container.

The concentration of eggs in the egg suspension was determined by counting the eggs using a microscope and a McMaster. The egg concentration was subsequently brought to a final concentration of 100 eggs per 200 μ L. To avoid proliferation of fungi, 5 μ g amphotericin B solution (Sigma, Germany) was added per millilitre of egg suspension.

Egg hatch assay

The in vitro EHA was based on the procedure described by Adamu et al. (2013), which is based on the method described by Coles et al. (1992). The egg suspension (200 µL) containing ~100 fresh eggs were distributed into each well of a 24-flatbottomed microtitre plate. The same volume (200 µL) of the dried plant extract (5 mg/mL) dissolved in solvent was added to each well. Albendazole was used as positive control, and the solvents were used as the negative controls. Albendazole was dissolved in 5% dimethyl sulphoxide (DMSO) in water and evaluated at various concentrations $(0.008 \ \mu g/mL - 25 \ \mu g/mL)$. The plates were incubated for 48 h at 27 °C at 70% relative humidity. The experiment was replicated three times for each extract on the same plate. After incubation, the hatched larvae and unhatched eggs were counted using an inverted microscope under 20 × magnification. The percentage inhibition of egg hatching was calculated using the formula of Bizimenyera et al. (2006):

TABLE 1: T	he plant and plant pai	rt used for the solvent extractior	i, plant family, solven	nt, the mass and percenta	ge yield of extract obtained.

Entry	Plant and plant part used in extraction	Plant family	Solvent	Mass of extract (g)	Yield of extract (%)
1	Aloe rupestris (leaves)	Asphodelaceae	Acetone	1.0127	5
2	Antizoma angustifolia (roots)	Menispermaceae	Acetone	1.0619	5
3	Calpurnia aurea (leaves, flowers)	Fabaceae	Acetone	0.9409	5
	Calpurnia aurea (stems)	Fabaceae	Acetone	1.0491	5
4	Senna italica subsp. arachoides (Burch.) Lock (roots, leaves, fruit)	Leguminosae	Acetone	1.0920	5
5	Cissus quadrangularis (stems)	Vitaceae	Acetone	1.0063	5
6	Clematis brachiata (whole plant)	Ranunculaceae	Acetone	1.0430	5
7	Cleome gynandra (leaves)	Capparidaceae	Acetone	0.9699	5
8	Ficus sycomorus (bark, stems)	Moraceae	Acetone	1.0339	5
9	Hypoxis rigidula (bulb)	Hypoxidaceae	Acetone	1.1049	6
10	Maerua angolensis (leaves)	Capparaceae	Acetone	1.1714	6
	Maerua angolensis (stem)	Capparaceae	Acetone	1.0524	5
11	Monsonia angustifolia (whole plant)	Geraniaceae	Acetone	1.0013	5
12	Pelargonium luridum (whole plant)	Geraniaceae	Acetone	1.0100	5
13	Schkuhria pinnata (whole plant)	Asteraceae	Acetone	1.1115	5
14	Sclerocarya birrea (bark, root)	Anacardiaceae	Acetone	0.9109	5
15	Sclerocarya birrea (fruit)	Anacardiaceae	Acetone	1.0142	5
16	Tabernaemontana elegans (leaves)	Apocynaceae	Acetone	1.0023	5

Egg hatch inhibition (%) = $100 (1 - \text{number of larvae})$	/
number of larvae and eggs in water control)	[Eqn 1]

The results of the EHA are shown in Table 2.

Determining the toxicity of the plant extracts

The toxicity of the plant extracts were determined by using the method employed by Adamu et al. (2013): Vero African Green monkey kidney cells were obtained from a confluent monolayer and then trypsinised and seeded (0.5×103 cells/ well) in a 96-well microtitre plate. This was followed by incubation overnight at 37 $^\circ\!C$ in 200 μL of 5% minimal essential medium (MEM, Highveld Biological, South Africa) and supplemented with 0.1% gentamicin (VirbacR) and 5% foetal calf serum (Adcock-Ingram) (Adamu et al. 2013). The media were replaced with 200 μ L of the extracts (1 mg/mL, 0.1 mg/mL, 0.01 mg/mL and 0.001 mg/mL) after 24 h and incubated for another 5 days. Viability of cells was determined using the tetrazolium-based colorimetric MTT assay (3-5-dimethyl thiazol-2-yl-2, 5-diphenyl tetrazolium bromide) as described by Mosmann (1983) (Adamu et al. 2013). Basically, the medium in each well was removed, replaced with fresh medium and $30 \,\mu\text{L} (5 \,\text{mg/mL})$ MTT in phosphate-buffered saline (PBS) followed by incubation for 4 h. The medium was then removed before washing the cells with PBS and before the addition of DMSO (50 µL) to dissolve any formazan crystals (Adamu et al. 2013). A Versamax microplate reader at 570 nm (path length 1 cm) was used to measure the absorbance of the wells. Doxorubicin was used as a positive control and tested at different concentrations. The negative control was a well containing cells without an extract. The percentage of cell viability was calculated relative to the pure growth. The LC_{50} value was calculated by determining the concentration of each plant extract resulting in 50% reduction of absorbance compared to untreated cells. Tests on the concentration of each extract were carried out in triplicate, and each experiment was repeated three times. The LC_{50} results are expressed as the mean \pm standard deviation (s.d.) of the three replicates. A plant extract having an LC_{50} value > 20 µg/mL has an acceptable level of toxicity, whilst a value < 20 µg/mL is regarded as toxic (Kuete & Efferth 2010).

Data analysis

Excel for Windows 7 was used to record the results produced in this study. Kinetica 5.0 (Thermo) using a sigmoid inhibitory model was used to calculate the LC_{50} values. The results are presented as the mean LC_{50} and the standard deviation of the mean.

Results

Extraction of plant material

Acetone was selected as extractant because it has many advantages over other generally used extractants (Eloff 1998). The extraction yield was very similar for the different plant species and plant parts and varied from 5% to 6% (Table 1).

Effect of different solvents on *Haemonchus* contortus egg hatching

The dried extracts did not dissolve in water, and therefore the use of acetone, DMSO and Tween 80 on the hatching of eggs. The final concentration of the solvents in the well was 50.0%, 25.0%, 12.5%, 6.3% and 3.1%. Even 3.1%, the lowest concentration of acetone, inhibited 94.0% egg hatching.

TABLE 2: Mean inhibition percentages of the acetone extracts (2.5 mg/mL) on egg hatching of *Haemonchus contortus* from sheep and the toxicity values (LC_{so}) against Vero cells.

Entry	Plant name and part used	Mean egg hatch inhibition (%) ± s.d.	Toxicity against Vero Cells LC ₅₀ (μg/mL) ± s.d.
1	Cleome gynandra (leaves)	68 ± 3	553.61 ± 18.83
2	Maerua angolensis (stem)	65 ± 5	180.64 ± 3.4
3	Monsonia angustifolia (whole plant)	56 ± 6	120.37 ± 4.06
4	Senna italica subsp. arachoides (Burch.) Lock (roots, leaves, fruit)	55 ± 13	46.31 ± 2.89
5	Aloe rupestris (leaves)	47 ± 7	63.46 ± 11.00
6	Tabernaemontana elegans (leaves)	47 ± 7	32.35 ± 0.88
7	Schkuhria pinnata (whole plant)	41 ± 14	39.93 ± 1.80
8	Antizoma angustifolia (roots)	37 ± 16	43.59 ± 6.28
9	Calpurnia aurea (stems)	32 ± 20	223.97 ± 5.3
10	Sclerocarya birrea (fruit)	28 ± 23	214.79 ± 14.0
11	Calpurnia aurea (leaves, flowers)	27 ± 16	166.63 ± 7.97
12	Pelargonium luridum (whole plant)	25 ± 10	30.58 ± 3.40
13	Maerua angolensis (leaves)	25 ± 6	73.76 ± 0.27
14	Ficus sycomorus (bark, stems)	25 ± 5	172.94 ± 8.91
15	Ficus sycomorus (stem)	21 ± 14	48.74 ± 1.32
16	Hypoxis rigidula (bulb)	17 ± 20	64.04 ± 2.53
17	Clematis brachiata (whole plant)	<i>11</i> ± <i>15</i>	117.00 ± 4.08
18	2.5% DMSO	7 ± 17	N.D.
		14 ± 13	N.D.
19	Water	16 ± 10	N.D.
		16 ± 9	N.D.
20	Doxorubicin	N.D.	2.97 ± 0.016

Note: Albendazole was the positive control and recorded 100% egg hatch inhibition at all concentrations (0.008 µg/mL – 25 µg/mL) used, whilst 2.5% DMSO recorded < 10% inhibition. N.D., not determined.

The lowest concentration of Tween 80 inhibited 19.0% egg hatching. DMSO concentrations of 25.0% and higher led to 100.0% inhibition of egg hatching. The best results were obtained with 3.1% and 6.3% DMSO leading to 13.0% and 19.0% inhibition of egg hatching, respectively. The results for the negative controls of water and PBS were 13.0% and 11.0%, indicating that 3.1% DMSO did not have a marked effect on the egg hatching. The lower degree of hatching with the negative controls probably indicates damage to the eggs during processing. Consequently, the extracts were dissolved in 6.0% DMSO leading to a 3.0% final concentration after adding the same volume of egg suspension.

Determining the inhibitory activity of the plant extracts on *Haemonchus contortus* egg hatching

The two negative controls also had a degree of activity on the egg hatching. The results are presented in Table 2 as the mean egg hatch inhibition (%) and the standard deviation of the mean. At 2.5 mg/mL, extracts of C. gynandra (leaves), M. angolensis (stem), M. angustifolia (whole plant) and S. italica subsp. arachoides (roots, leaves and fruit) had a mean inhibition rate of between 55% and 68% which was much higher than the water and DMSO negative controls. Our choice of 2.5 mg/mL appeared to be a good concentration because only four species had what would have been an LC_{50} in the order of 2.5 mg/mL. Albendazole, the positive control, recorded 100% inhibition at the lowest concentration, 0.008 µg/mL. The extract of C. brachiata (whole plant) had the lowest activity (11%), which was even lower than the water control group (16%). The activity of the plant extracts was so much lower than that of albendazole that the feasibility of using plant extracts can be questioned. Plant extracts may be more active in the larval development assay (LDA).

Determining the cytotoxicity of the plant extracts

The tetrazolium-based (MTT) colorimetric assay (Mosmann 1983) was used to determine the viability of Vero African Green monkey kidney cells in the presence of each of the plant extracts and the results are shown in Table 2.

From these results, it is apparent that none of the plant acetone extracts were as toxic as doxorubicin (2.97 μ g/mL ± 0.016 μ g/mL = 5.12 μ M ± 0.028 μ M). The leaf extract of *C. gynandra* was the least toxic (LC₅₀ = 553.61 μ g/mL ± 18.83 μ g/mL) followed by the stem extract of *C. aurea* (LC₅₀ = 223.97 μ g/mL ± 5.4 μ g/mL), the fruit extract of *S. birrea* (LC₅₀ = 214.79 μ g/mL ± 14 μ g/mL), the stem extract of *M. angolensis* (LC₅₀ = 180.64 μ g/mL ± 3.5 μ g/mL) and the bark and stem extract of *F. sycomorus* (LC₅₀ = 172.94 μ g/mL ± 8.91 μ g/mL). The whole plant extract of *P. luridum* (LC₅₀ = 30.58 μ g/mL ± 3.40 μ g/mL) was the most toxic of all the plants against Vero cells.

Discussion

The EHA is an *in vitro* assay used to evaluate the anthelmintic activities of natural products. The capacity to reduce egg

hatching could help to modulate the risk of parasitism by limiting the infectivity of pastures grazed by ruminants (Max 2010).

The aim of this study was to determine the inhibitory activity of the acetone extracts of 15 plant species on egg hatching of H. contortus in order to select the most promising plant species that could control the nematodes in the animal gut for further study. In previous studies, it was found that aqueous extracts contained few compounds, had very low biological activity (Eloff, Famakin & Katerere 2005; Kotze & Eloff 2002) and had low or negligible anthelmintic activity (Bizimenyera et al. 2006; Worku, Franco & Miller 2009). Acetone was therefore selected as an appropriate extractant because it is miscible with organic and aqueous solvents, non-toxic to bacteria and fungi, and also has the capacity to extract a wide range of polar compounds (Eloff 1998). As shown in Table 2, it is evident that the extracts of four plant species had anthelmintic activity (inhibitory activity above 50%) at the concentration tested.

Cleome gynandra leaf extracts had the best anthelmintic activity with an egg hatch inhibition of $68\% \pm 3\%$ and low toxicity (LC₅₀ = 553.61 μ g/mL ± 18.83 μ g/mL) on Vero cells. Our results are in agreement with that of other researchers who have also reported on the anthelmintic activity of C. gynandra. Two authors (Jadhav, Ghawate & Bhamber 2011; Thenmozhi et al. 2014) used the unverified assumption that the Indian adult earthworm (Pheretima posthuma) could be used as a model for the activity of C. gynandra (syn. Gynandropsis pentaphylla) extracts against intestinal roundworm parasites of human beings because of its anatomical and physiological resemblance. They also used a physiologically non-relevant high concentration of 25 mg/mL and concluded that these extracts had potent anthelmintic activity when it killed earthworms after 53 min without examining any helminths. Sowunmi and Afolayan (2015) also did a phytochemical analysis of the acetone extract of different parts of C. gynandra. The polyphenolic contents of the various parts of the plant were significantly high. Leaf acetone extracts of C. gynandra had the highest concentration of total phenolics (126.79 mg/g \pm 0.55 mg/g), flavonoids (40.58 mg/g \pm 0.06 mg/g) and flavanols (42.41 mg/g \pm 0.05 mg/g), whilst the stem extract had the highest amount of proanthocyanidins (419.01 mg/g \pm 0.67 mg/g) compared to the leaves (403.29 mg/g \pm 0.89 mg/g) and fruits (107.18 mg/g \pm 0.59 mg/g). The low concentration of saponins and alkaloids suggests that this plant may have low toxicity (Sowunmi & Afolayan 2015). This suggestion is supported by the low toxicity we observed against Vero cells (LC₅₀ = $553.61 \,\mu g/mL \pm 18.83 \,\mu g/mL$).

Alcohol and aqueous extracts from the leaves of *Cleome viscosa* Linn were also investigated for their anthelmintic activity against the adult Indian earthworm, *P. posthuma*, as well as *Ascaridia galli*. Three concentrations (50 mg/mL, 100 mg/mL and 150 mg/mL) of each extract were studied,

which entailed the determination of time of paralysis and time of death of the worm. Both the extracts had significant anthelmintic activity at the highest concentration of 150 mg/mL. The water leaf extract had weaker activity than the methanolic leaf extract, and both extracts caused paralysis and death of worms. Phytochemical screening of the methanol extract showed that anthraquinone glycosides, phenolic compounds and steroids were present in C. viscosa Linn, whilst in the aqueous extract glycosides and phenolic compounds were present. Flavonoids were identified as being one of the chemical constituents amongst the phenolic compounds in the crude extracts. Polyphenolic compounds are known for their anthelmintic activity (Kaushik, Katiyar & Sen 1974; Lal et al. 1976; Szewezuk, Mongelli & Pomilio 2003). Synthetic phenolic anthelmintics such as niclosamide, oxyclozanide and bithionol interfere with energy generation in helminth parasites by uncoupling oxidative phosphorylation (Bate-Smith 1962; Martin 1997; Tandon et al. 1997). The phenolic content may therefore have produced similar activity in the extracts of C. viscosa Linn and C. gynandra.

The second best egg hatch inhibition of $65\% \pm 5\%$ was by *M. angolensis* (stem) extract that had low toxicity ($LC_{50} = 180.64$ μ g/mL ± 3.5 μ g/mL) on Vero cells. Phytochemical screening of the methanolic extract of the stem bark found that glycosides, tannins, saponins, terpenes, flavonoids, carbohydrates, proteins and alkaloids were present in M. angolensis (Ayo et al. 2013; Meda et al. 2013; Pl@ntUse). These compounds could also be present in the acetone stem extract tested in this study. It has been reported that Maerua edulis (Gilg and Gilg-Ben.) DeWolf and Maerua subcordata (Gilg) DeWolf have been used in traditional anthelmintic remedies in Kenya to treat sheep infected with H. contortus (Gakuya 2001). In this study, aqueous extract from both unground and ground material of each plant material was prepared using boiling water. Twenty-one clinically healthy sheep of mixed breeds and sexes were randomly allocated to four treatment groups, four animals each. Faecal egg counts were performed for all the sheep. It was found that the crude extracts could control helminthoses to a reasonable extent and maintain the animal at a clinically healthy state. The Brine shrimp assay was used to detect bioactivity in the water, chloroform and methanol extracts of M. subcordata and M. edulis. The chloroform extract was the most toxic to the Brine shrimps compared to the water and methanol extracts (Gakuya 2001).

Monsonia angustifolia (whole plant) had an egg hatch inhibition of $56\% \pm 6\%$ and also had low toxicity on Vero cells (LC₅₀ = 120.37 µg/mL ± 4.06 µg/mL). Five compounds identified as aryl naphthalene lignans (5-methoxyjusticidin A, justicidin A, chinensinaphthol, retrochinensinaphthol methyl ether and suchilactone) were isolated during the fractionation of the organic (methanol-dichloromethane) extract of *M. angustifolia* (Khorombi 2006). Lignans are a group of naturally occurring phenolic compounds. The drug podophyllum, a lignan, is obtained from the dried root and rhizomes of two species of *Podophyllum* (Berberidaceae), the American species *Podophyllum peltatum* and the Indian species *Podophyllum hexandrum* (*Podophyllum emodi*). The European settlers reported using the root extensively, particularly as a cathartic and anthelmintic (Konuklugil 1995). Thus, the anthelmintic activity of *M. angustifolia* may be attributed to the lignans.

Senna italica subsp. arachoides (Burch.) Lock (roots, leaves, fruit) had an egg hatch inhibition of 55% ± 13% and had a higher toxicity on Vero cells ($LC_{50} = 46.31 \,\mu\text{g/mL} \pm 2.89 \,\mu\text{g/mL}$) than that of *C. gynandra* (leaves), *M. angolensis* (stem) and *M. angustifolia* (whole plant) which also had anthelmintic activity.

Aloe rupestris (leaves) only had an egg inhibition of $47\% \pm 7\%$ against *H. contortus* and low toxicity (LC₅₀ = 63.46 ± 11.00) against Vero cells. It has been reported that other Aloe species such as *Aloe ferox* can affect *H. contortus* of goats negatively (Maphosa *et al.* 2010). The amino acids, saponins and sterols in *A. ferox* can disturb protein structure and therefore affect the growth and repair of the nematode body (Mabusela, Stephen & Botha 1990).

Tabernaemontana elegans (leaves) also only had an egg inhibition of 47% \pm 7% and low toxicity (LC₅₀ = 32.35 \pm 0.88) against Vero cells. In Guadeloupe (French West Indies), another species, Tabernaemontana citrifolia, has traditionally been used as an anthelmintic preparation for ruminants (Marie-Magdeleine et al. 2010). Marie-Magdeleine et al. (2010) prepared aqueous, methanolic and dichloromethane extracts from the fruit, leaves and roots of T. citrifolia for testing on four developmental stages of *H. contortus*. The EHA, the LDA, the L3 migration inhibition assay (LMI) and the adult worm motility assay (AWM) were employed in the testing. From the tests it was apparent that there were significant effects for the different parts of T. citrifolia when compared to the negative control and that the differences depended on the parasitic stage. The efficacies on the larval development of H. contortus ranged from 88.9% to 99.8% for fruit, from 72.1% to 83.8% for roots and from 33.5% to 85.0% for leaves. For the methanolic extract a dose-dependent effect was observed. Alkaloid compounds are present in the different parts of T. citrifolia and may be responsible for the observed anthelmintic activity against H. contortus (Marie-Magdeleine et al. 2010). Another species, Tabernaemontana coronaria, was also investigated for anthelmintic activity against the Indian adult earthworm P. posthuma. It was found that the ethanolic extract of the leaves had potent anthelmintic activity (Pushpa et al. 2011).

Anthelmintic activity has also been documented for other plant species where an organic extract was also used in the determination of the anthelmintic activity (Ademola & Eloff 2010, 2011; Monteiro *et al.* 2010). Adamu *et al.* (2013) reported that *Heteromorpha trifoliate, Leucosidia sericea* and *Maesa lanceolata* had 100% inhibition, whilst *Clausenia anisata* had 80% inhibition at a concentration of 3.13 mg/mL in the EHA. Several other plant species have been documented as having anthelmintic activity. These are *Lantana camara* (Verbenaceae), *Tagetes minuta* (Asteraceae), *Mentha villosa* (Lamiaceae) (Albuquerque *et al.* 2007) and *Alpinia zerumbet* (Zingiberaceae) (Almeida 1993). In this study, extracts of *C. gynandra* (leaves), *M. angolensis* (stem), *M. angustifolia* (whole plant) and *S. italica* subsp. *arachoides* (roots, leaves and fruit) only had inhibition activity between 55% and 68% at 2.5 mg/mL. Higher inhibitory activity for these extracts may have been obtained if they had also been evaluated at 3.13 mg/mL.

The kidney is one of the main sites of excretion in animals, and therefore, renal cells in culture were used as an indicator of toxicity for this study. These cells were explicitly chosen because of the favoured blood supply of the kidney and their high metabolic capacity. The results of the toxicity study were encouraging because the extracts had low toxicity ($LC_{50} > 20 \ \mu g/mL$) against Vero cells. The determination of cellular toxicity is valuable because it will give a good indication of whether *in vitro* toxicity is also an indicator of *in vivo* toxicity.

The results of this study are significant because egg hatch inhibition is an important method by which pasture contamination by animals can be reduced during grazing. Thus, there is the possibility that the plant material in this study can be administered as a feed to control helminths, although the concentrations required for efficacy were very high. Overall, the use of these botanicals could contribute to a helminth control programme.

Conclusion

The acetone extracts of *C. gynandra*, *M. angolensis*, *M. angustifolia* and *S. italica* subsp. *arachoides* have some anthelmintic activity against *H. contortus* egg hatching. The activity of each of these extracts was lower than that of the positive control, albendazole. The most promising plant species is *C. gynandra*, which may be further studied to identify the active constituents responsible for anthelmintic activity.

Acknowledgements

The authors thank the Technology Innovation Agency (TIA) of South Africa for financial support and Professor Lyndy McGaw for informative discussions about the toxicity of plant extracts on Vero cells.

Competing interests

The authors declare that they have no financial or personal relationships that may have inappropriately influenced them in writing this article.

Authors' contributions

G.F. conceptualised the study. G.F., K.W.W. and T.L. did the literature search and plant selection. T.L. prepared the plant extracts. J.N.E. and E.P. arranged for sheep to be infected, collected the eggs and guided the study. B.M.S. did the egg hatch assay on the extracts. O.T.A. screened the extracts for toxicity on Vero cells. K.W.W. wrote the first draft of the manuscript.

References

- Adamu, M., Naidoo, V. & Eloff, J.N., 2013, 'Efficacy and toxicity of thirteen plant leaf acetone extracts used in ethnoveterinary medicine in South Africa on egg hatching and larval development of *Haemonchus contortus*', *BMC Veterinary Research* 9, 38. http://dx.doi.org/10.1186/1746-6148-9-38
- Adejinmi, J.O. & Harrison, L.J.S., 1997, 'Parasitic nematodes of domestic ruminants in Nigeria: Impact on ruminant production and control', *Tropical Veterinarian* 15, 137–148.
- Ademola, I.O. & Eloff, J.N., 2010, 'In vitro anthelmintic activity of *Combretum molle* (R. Br. ex G. Don) (Combretaceae) against *Haemonchus contortus* ova and larvae', *Veterinary Parasitology* 169, 198–203. http://dx.doi.org/10.1016/j.vetpar.2009. 12.036
- Ademola, I.O. & Eloff, J.N., 2011, 'Anthelmintic activity of acetone extract and fractions of Vernonia amygdalina against Haemonchus contortus eggs and larvae', Tropical Animal Health and Production 43, 521–527. http://dx.doi.org/10.1007/s11250-010-9727-7
- Akhtar, M.S., Igbal, Z., Khan, M.N. & Lateef, M., 2000, 'Anthelmintic activity of medicinal plants with particular reference to their use in animals in the Indo– Pakistan subcontinent', *Small Ruminants Research* 38, 99–107. http://dx.doi. org/10.1016/S0921-4488(00)00163-2
- Albuquerque, U.P., Medeiros, P.M., Almeida, A.L., Monteiro, J.M., Neto, E.M.F.L., Melo, J.G. et al., 2007, 'Medicinal plants of the caatinga (semi-arid) vegetation of NE Brazil: A quantitative approach', *Journal of Ethnopharmacology* 114, 325–354. http://dx.doi.org/10.1016/j.jep.2007.08.017
- Almeida, E.R. (ed.), 1993, Plantas medicinais brasileiras conhecimentos populares e científicos, Hemus, São Paulo, p. 341.
- Angulo-Cubillán, F.J., García-Coiradas, L., Alunda, J.M., Cuquerella, M. & De La Fuente, C., 2010, 'Biological characterization and pathogenicity of three *Haemonchus contortus* isolates in primary infections in lambs', *Veterinary Parasitology* 171, 99–105. http://dx.doi.org/10.1016/j.vetpar.2010.03.004
- Arosemena, N.A.E., Bevilaqua, C.M.L., Melo, A.C.F.L. & Girão, M.D., 1999, 'Seasonal variations of gastrointestinal nematodes in sheep and goats from semi-arid area in Brazil', *Revista Medicina Veterinária* 150, 873–876.
- Ayo, R.G., Audu, O.T., Amupitan, J.O. & Uwaiya, E., 2013, 'Phytochemical screening and antimicrobial activity of three plants used in traditional medicine in Northern Nigeria', Journal of Medicinal Plants Research 7(5), 191–197.
- Bate-Smith, E.C., 1962, 'The phenolic constituent of plants and their taxonomic significance, dicotyledons', *Journal of the Linnean Society, Botany* 58, 95–103.
- Batista, L.M., Bevilaqua, C.M.L., Moraes, S.M. & Vieira, L.S., 1999, 'In vitro ovicidal and larvicidal effect of the plants Spigelia anthelmia and Momordica charantia against the nematode Haemonchus contortus', Ciência Animal 9, 67–73.
- Bizimenyera, E.S., Githiori, J.B., Eloff, J.N. & Swan, G.E., 2006, 'In vitro activity of Peltophorum africanum Sond. (Fabaceae) extracts on the egg hatching and larval development of the parasitic nematode Trichostrongylus colubriformis', Veterinary Parasitology 142, 336–343. http://dx.doi.org/10.1016/j.vetpar.2006.06.013
- Coles, G.C., Bauer, C., Borgsteede, F.H.M., Geerts, S., Klei, T.R., Taylor, M.A. et al., 1992, 'World Association for the Advancement of Veterinary Parasitology (WAAVP) methods for the detection of anthelmintic resistance in nematodes of veterinary importance', Veterinary Parasitology 44, 35–44. http://dx.doi.org/10.1016/0304-4017(92)90141-U
- Eloff, J.N., 1998, 'Which extractant should be used for the screening and isolation of antimicrobial components from plants?', *Journal of Ethnopharmacology* 60, 1–8. http://dx.doi.org/10.1016/S0378-8741(97)00123-2
- Eloff, J.N., Famakin, J.O. & Katerere, D.R.P., 2005, 'Combretum woodii (Combretaceae) leaf extracts have high activity against Gram-negative and Gram-positive bacteria', African Journal of Biotechnology 4, 1161–1166.
- Gakuya, D.W., 2001, 'Pharmacological and clinical evaluation of anthelmintic activity of Albizia anthelmintica Brogn, Maerua edulis De Wolf and Maerua subcordata De Wolf plant extracts in sheep and mice', PhD., University of Nairobi, Department of Veterinary Clinical Studies.
- Hammond, J.A., Fielding, D. & Bishop, S.C., 1997, 'Prospects for plant anthelmintics in tropical veterinary medicine', Veterinary Research Communications 21, 213–228. http://dx.doi.org/10.1023/A:1005884429253
- Hounzangbe-Adote, M.S., Paolini, V., Fouraste, I., Moutairou, K. & Hoste, H., 2005, 'In vitro effects of four tropical plants on three life-cycle stages of the parasitic nematode, *Haemonchus contortus'*, *Research in Veterinary Science* 78, 155–160. http://dx.doi.org/10.1016/j.rvsc.2004.05.009
- Jadhav, V.S., Ghawate, V.B. & Bhamber, R.S., 2011 'Anthelmintic activity of ethanolic extract of seeds of Cleome gynandra Linn', Journal of Pharmacy Research 4(12), 4479.
- Kaushik, R.K., Katiyar, J.C. & Sen, A.B., 1974, 'Studies on the mode of the action of anthelmintics with Ascaridia galli as a test parasite', Indian Journal of Medical Research 62, 1367–1375.
- Khorombi, T.E., 2006, 'A chemical and pharmacological investigation of three South African plants', M.Sc dissertation, University of KwaZulu Natal South Africa, Pietermaritzburg, School of Chemistry.
- Kuete, V. & Efferth, T., 2010, 'Cameroonian medicinal plants: Pharmacology and derived natural products', Frontiers in Pharmacology 1(123), 1–19. http://dx.doi. org/10.3389/fphar.2010.00123
- Konuklugil, B., 1995, 'The importance of aryltetralin (podophyllum) lignans and their distribution in the plant kingdom', Journal of Faculty of Pharmacy of Ankara 24, 2.
- Kotze, M. & Eloff J.N., 2002, 'Extraction of antibacterial compounds from Combretum microphyllum (Combretaceae)', South African Journal of Botany 68, 62–67. http:// dx.doi.org/10.1016/S0254-6299(16)30456-2

- Lal, J., Chandra, S., Raviprakash, V. & Sabir, M., 1976, 'In vitro anthelmintic action of some indigenous medicinal plants on Ascaridia galli worms', Indian Journal of Physiology and Pharmacology 20, 64–68.
- Mabusela, W.T., Stephen, A.M. & Botha, M.C., 1990, 'Carbohydrate polymers from Aloe ferox leaves', Phytochemistry 29, 3555–3558. http://dx.doi.org/10.1016/ 0031-9422(90)85275-K
- Maphosa, V., Masika, P.J., Bizimenyera, E.S. & Eloff J.N., 2010. 'In vitro anthelmintic activity of crude aqueous extracts of Aloe ferox, Leonotis leonurus and Elephantorrhiza elephantina against Haemonchus contortus', Tropical Animal Health and Production 42, 301–307. http://dx.doi.org/10.1007/s11250-009-9421-9
- Marie-Magdeleine, C., Mahieu,, M., D'Alexis, S., Philibert, L. & Archimede, H., 2010, 'In vitro effects of Tabernaemontana citrifolia extracts on Haemonchus contortus', Research in Veterinary Science 89(1), 88–92. http://dx.doi.org/10.1016/j.rvsc.2010.01.002
- Martin, R.J., 1997, 'Mode of action of anthelmintic drugs', Veterinary Journal 154, 11–34. http://dx.doi.org/10.1016/S1090-0233(05)80005-X
- Max, R.A., 2010. 'Effect of repeated wattle tannin drenches on worm burdens, faecal egg counts and egg hatchability during naturally acquired nematode infections in sheep and goats', Veterinary Parasitology 169(1–2), 138–143. http://dx.doi. org/10.1016/j.vetpar.2009.12.022
- Meda, N.T.R., Bangou, M.J., Bakasso, S., Millogo-Rasolodimby, J. & Nacoulma, O.G., 2013, 'Antioxidant activity of phenolic and flavonoid fractions of *Cleome gynandra* and *Maerua* angolensis of Burkina Faso', *Journal of Applied Pharmaceutical Science* 3(02), 036–042.
- Melo, A.C.F.L., Bevilaqua, C.M.L. & Reis, I.F., 2009, 'Anthelmintic resistance to benzimidazole in gastrointestinal nematodes from small ruminants of semi-arid Brazilian northeast', *Ciência Animal* 10, 294–300.
- Mendoza de Gives, P., Crespo, J.F., Rodriguez, D.H., Prats, V.V., Hernandez, E.L. & Fernandez, G.E.O., 1998, 'Biological control of *Haemonchus contortus* infective larvae in ovine faeces by administering an oral suspension of *Duddingtonia flagrans* chlamydospores to sheep', *Journal of Helminthology* 72, 343–347. http:// dx.doi.org/10.1017/S0022149X00016710
- Monteiro, M.V.B., Bevilaqua, C.M.L., Morais, S.M., Machado, L.K.A., Camurca-Vasconcelos, A.L.F., Ampello, C.C. et al., 2010, 'Anthelmintic activity of Jatropha curcas L. seeds on Haemonchus Contortus', Veterinary Parasitology 182, 259–263. http://dx.doi.org/10.1016/j.vetpar.2011.04.010
- Mosmann, T., 1983, 'Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays', *Journal of Immunological Methods Methods* 65(1–2), 55–63. http://dx.doi.org/10.1016/0022-1759(83)90303-4

- Pl@ntUse, viewed 16 December 2015, from http://uses.plantnet-project.org/en/ Maerua_angolensis_(PROTA)
- Prichard, R., 1994, 'Anthelmintic resistance', Veterinary Parasitology 54, 259–268. http://dx.doi.org/10.1016/0304-4017(94)90094-9
- Pushpa, B., Latha, K.P., Vaidya, V.P., Shruthi, A. & Shweath, C., 2011, 'In vitro anthelmintic activity of leaves extracts of *Tabernaemontana coronaria*', *International Journal of ChemTech Research* 3(4), 1788–1790.
- Schnyder, M., Torgerson, P.R., Schonmann, M., Kolera, L. & Hertzberg, H., 2005, 'Multiple anthelmentic resistance in *Haemonchus contortus* isolated from South African Boer goats in Switzerland', *Veterinary Parasitology* 128, 285–290. http:// dx.doi.org/10.1016/j.vetpar.2004.12.010
- Slomp, L., Pereira, P.S., Zingaretti, S.D. & Beleboni, R.O., 2009, *In vitro* nematocidal effects of medicinal plants from the Sao Paulo state-Brazil', *Pharmaceutical Biology* 47, 230–235.
- Sowunmi, L.I. & Afolayan, A.J., 2015, 'Phytochemical constituents and antioxidant properties of acetone extract of *Cleome gynandra* (L.) growing in the Eastern Cape, South Africa', *African Journal of Traditional Complementary and Alternative Medicines* 12(3), 1–8. http://dx.doi.org/10.4314/ajtcam.v12i3.1
- Szewezuk, V.D., Mongelli, E.R. & Pomilio, A.B., 2003, 'Antiparasitic activity of Melia azadirach growing in Argentina', Molecular Medicinal Chemistry 1, 54–57.
- Tandon, V., Pal, P., Roy, H.S. & Reddy, K.S., 1997, 'In vitro anthelmintic activity of root-tuber extract of Flemingia vestita, an indigenous plant in ShillongIndia', *Parasitology Research* 83, 492–498. http://dx.doi.org/10.1007/ s004360050286
- Thenmozhi, S., Varghese, N., Subasini, U., Dhanalakshmi, M., Manjuladevi, K. & Kameshwaran, S., 2014 'Comparative study of anthelmintic activity of different extracts of Gynandropsis pentaphyla Linn', International Journal of Pharmacy and Life Sciences 5(1), 3246–3248.
- Torres-Acosta, J.F.J., Mendoza-de-Gives, P., Aguilar-Caballero, A.J. & Cuellar-Ordaz, J.A., 2012, 'Anthelmintic resistance in sheep farms: Update of the situation in the American continent', Veterinary Parasitology 189, 89–96. http://dx.doi. org/10.1016/j.vetpar.2012.03.037
- Waller, P.J., 1994, 'The development of anthelmintic resistance in ruminant livestock', Acta Tropica 56, 233–243. http://dx.doi.org/10.1016/0001-706X(94)90065-5
- Worku, M., Franco, R. & Miller, J.H., 2009, 'Evaluation of the activity of plant extracts in Boer goats', American Journal of Animal and Veterinary Sciences 4, 2–79. http:// dx.doi.org/10.3844/ajavsp.2009.72.79