



Molecular biological characteristics of foot-and-mouth disease virus in the African buffalo in southern Africa

Authors:

Christopher J. Kasanga¹
 Rahana Dwarka²
 Gaothlele Thobokwe³
 Jemma Wadsworth⁴
 Nick J. Knowles⁴
 Misheck Mulumba⁵
 Ezekia Ranga⁶
 Jimis Deve^{1,11}
 Cornelius Mundia¹⁰
 Patrick Chikungwa⁸
 Laureta Joao⁹
 Raphael Sallu⁷
 Mmeta Yongolo⁷
 Philemon N. Wambura¹
 Mark M. Rweyemamu¹
 Donald P. King⁴

Affiliations:

¹Southern African Centre for Infectious Diseases Surveillance, Sokoine University of Agriculture, Tanzania

²Onderstepoort Veterinary Institute, University of Pretoria, South Africa

³Botswana Vaccine Institute, Botswana

⁴WRLFMD, The Pirbright Institute, United Kingdom

⁵Southern African Development Community Secretariat, Botswana

⁶Ministry of Livestock Development and Fisheries, Tanzania

⁷Tanzania Veterinary Laboratory Agency, Tanzania

⁸Southern African Development Community, Transboundary Animal Diseases Section, Malawi

⁹Southern African Development Community, Transboundary Animal Diseases Section, Angola

¹⁰Southern African Development Community, Transboundary Animal Diseases Section, Zambia

¹¹Southern African Development Community, Transboundary Animal Diseases Section, Mozambique

Read online:



Scan this QR code with your smart phone or mobile device to read online.

Foot-and-mouth disease (FMD) is endemic in most countries in southern Africa. African buffaloes (*Syncerus caffer*) are known to play a significant role in the transmission and dynamics of FMD virus (FMDV) in wildlife-livestock interface areas. The aim of this study was to investigate the serotype and determine the genetic relationships of FMDV recovered from animals in Tanzania, Zambia and Mozambique, and compare them with viruses detected from elsewhere in the sub-Saharan region. A total of 150 oesophageal-pharyngeal (probang) samples collected in 2010 from cattle and buffaloes in Marromeu (Mozambique), Katavi (Tanzania) and Lochinvar (Zambia) National Parks were used in this study. The presence of FMDV was determined by laboratory methods such as VI, antigen ELISA and real-time RT-PCR. Phylogenies of VP1 sequences were determined by the neighbour-joining method. The overall FMDV genome detection rate was 6.7% ($n = 10$), with SAT 1 being the most frequent serotype (60%; $n = 6$) isolated in cattle and buffaloes in Mozambique, Tanzania and Zambia followed by SAT 3 (30%; $n = 3$) and SAT 2 (10%; $n = 1$). Genotyping showed that type SAT 1 viruses fell into either the TOPOTYPE 1 (NWZ) or UNASSIGNED topotypes, type SAT 2 into the AFRICA topotype I and type SAT 3s into topotype IV (SEZ). This study reveals that serotypes SAT 1–3 are maintained in cattle and buffaloes in livestock-wildlife interface areas in Marromeu, Katavi, and Lochinvar National Parks. Phylogenetic analysis of FMDV isolates from Tanzania, Zambia and Mozambique showed that they are genetically related to lineages and topotypes from Africa. This information contributes to the understanding of the epidemiology of FMD in southern Africa. In Mozambique, Tanzania and Zambia, lack of consistent surveillance systems and animal movement controls make it difficult to determine the exact source of FMD and transmission dynamics of FMDV. Further studies are needed to elucidate the complex epidemiology of FMD in Africa.

Acknowledgements

This work was supported by the Wellcome Trust Grant WT087546MA to the Southern African Centre for Infectious Diseases & Surveillance (SACIDS).

Correspondence to: Christopher Kasanga

Email: christopher.kasanga@sacids.org

Postal address: PO Box 3019, Chuo Kikuu, Morogoro, Tanzania

How to cite this article: Kasanga, C.J., Dwarka, R., Thobokwe, G., Wadsworth, J., Knowles, N.J., Mulumba, M. *et al.*, 2014, 'Molecular biological characteristics of foot-and-mouth disease virus in the African buffalo in southern Africa', *Onderstepoort Journal of Veterinary Research* 81(2), Art. #728, 1 page. <http://dx.doi.org/10.4102/ojvr.v81i2.728>

Note: Proceedings of the 2nd One Health Conference in Africa. Jointly organised by the Southern African Centre for Infectious Disease Surveillance and the Tanzania National Institute for Medical Research, held at the Snow Crest Hotel in Arusha, Tanzania from 16th to 19th April 2013: <http://www.sacids.org/kms/frontend/index.php?m=119>.

Copyright: © 2014. The Authors. Licensee: AOSIS OpenJournals. This work is licensed under the Creative Commons Attribution License.